

PACKAGE LEAFLET: INFORMATION FOR THE USER

CENOMAR

400mg film-coated tablets

MOXIFLOXACIN

• This leaflet is a copy of the Summary of Product Characteristics and Patient Information Leaflet for a medicine, which outlines the conditions under which the medicine should be used and information on its known safety • The product information may be updated several times within its shelf life, and there could be differences between the version of information shown here and other information in the public domain or in the package insert • This leaflet may not contain all the information about the medicine or the information may not be the most up to date version for this product • If you have any questions or are not sure about anything, ask your doctor or pharmacist • Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

• Keep this leaflet • You may need to read it again • If you have any further questions, ask your doctor or pharmacist • This medicine has been prescribed for you only • Do not pass it on to others • It may harm them, even if their signs of illness are the same as yours • If you get any side effects, talk to your doctor or pharmacist • This includes any possible side effects not listed in this leaflet •

What is in this leaflet?

1. What Cenomar is and what it is used for
2. What you need to know before you take Cenomar
3. How to take Cenomar
4. Possible side effects
5. How to store Cenomar

1. WHAT CENOMAR IS AND WHAT IT IS USED FOR

Cenomar contains the active substance moxifloxacin, which belongs to a group of antibiotics called fluoroquinolones. Cenomar works by killing bacteria that cause infections.

Cenomar is used in patients aged 18 years and above for treating the following bacterial infections when caused by bacteria against which moxifloxacin is active. Cenomar should only be used to treat these infections when usual antibiotics cannot be used or have not worked:

Infection of the sinuses, sudden worsening of long term inflammation of the airways or infection of the lungs (pneumonia) acquired outside the hospital (except severe cases).

Mild to moderate infections of the female upper genital tract (pelvic inflammatory disease), including infections of the fallopian tubes and infections of the uterus mucous membrane.

Cenomar tablets are not sufficient on their own for treating this kind of infection. Therefore, another antibiotic in addition to Cenomar tablets should be prescribed by your doctor for the treatment of infections of the female upper genital tract (see section 2. What you need to know before you take Cenomar, Warnings and precautions, Talk to your doctor before taking Cenomar).

If the following bacterial infections have shown improvement during initial treatment with Cenomar solution for infusion, Cenomar tablets may also be prescribed by your doctor to complete the course of therapy:

Infection of the lungs (pneumonia) acquired outside the hospital, infections of the skin and soft tissue.

Cenomar tablets should not be used to initiate therapy for any type of infections of the skin and soft tissue or in severe infections of the lungs.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE CENOMAR

Contact your doctor if you are not sure if you belong to a patient group described below.

Do not take Cenomar

- If you are allergic to the active substance moxifloxacin, any other quinolone antibiotics or any of the other ingredients of this medicine.
- If you are pregnant or are breast-feeding.
- If you are under 18 years of age.
- If you have previously had problems with your tendons related to treatment with quinolone antibiotics (see section Warnings and Precautions ... and section 4. Possible side effects).
- If you were born with or have
 - any condition with abnormal heart rhythm (seen on ECG, electrical recording of the heart)
 - a salt imbalance in the blood (especially low levels of potassium or magnesium in the blood)
 - a very slow heart rhythm (called 'bradycardia')
 - a weak heart (heart failure)
 - a history of abnormal heart rhythmsor
 - if you are taking other medicines that result in abnormal ECG changes (see section Other medicines and Cenomar). This is because Cenomar can cause changes on the ECG, that is a prolongation of the QT-interval, i.e., delayed conduction of electrical signals.
- If you have a severe liver disease or increased liver enzymes (transaminases) higher than 5 times the upper normal limit.

Warnings and precautions

Talk to your doctor before taking Cenomar

- Cenomar can change your heart's ECG, especially if you are female, or if you are elderly. If you are currently taking any medicine that decreases your blood potassium levels, consult your doctor before taking Cenomar (see also sections Do not take and Other medicines and Cenomar).
- If you suffer from epilepsy or a condition which makes you likely to have convulsions talk to your doctor before taking Cenomar.
- If you have or have ever had any mental health problems, consult your doctor before taking Cenomar.
- If you suffer from myasthenia gravis (abnormal muscle fatigue leading to weakness and in serious cases paralysis), taking Cenomar may worsen the symptoms of your disease. If you think you are affected consult your doctor immediately.
- If you or any member of your family have glucose-6-phosphate dehydrogenase deficiency (a rare hereditary disease), tell your doctor, who will advise whether Cenomar is suitable for you.
- If you have a complicated infection of the female upper genital tract (e.g. associated with an abscess of the fallopian tubes and ovaries or of the pelvis), for which your doctor considers an intravenous treatment necessary, treatment with Cenomar tablets is not appropriate.
- For the treatment of mild to moderate infections of the female upper genital tract your doctor should prescribe another antibiotic in addition to Cenomar. If there is no improvement in symptoms after 3 days of treatment, please consult your doctor.

When taking Cenomar

- If you experience palpitations or irregular heart beat during the period of treatment, you should inform your doctor immediately. He/she may wish to perform an ECG to measure your heart rhythm.
- The risk of heart problems may increase with increase of the dose. Therefore, the recommended dosage should be followed.
- There is a rare chance that you may experience a severe, sudden allergic reaction (an anaphylactic reaction/shock) even with the first dose. Symptoms include tightness in the chest, feeling dizzy, feeling

sick or faint, or dizziness when standing up. If so, stop taking Cenomar and seek medical advice immediately.

- Cenomar may cause a rapid and severe inflammation of the liver which could lead to life-threatening liver failure (including fatal cases, see section 4. Possible side effects). If you suddenly feel unwell and/or are being sick and also have yellowing of the whites of the eyes, dark urine, itching of the skin, a tendency to bleed or liver induced disease of the brain (symptoms of a reduced liver function or a rapid and severe inflammation of the liver) please contact your doctor before taking any more tablets.
- If you develop a skin reaction or blistering / peeling of the skin and/or mucosal reactions (see section 4. Possible side effects) contact your doctor immediately before you continue treatment.
- Quinolone antibiotics, including Cenomar, may cause convulsions. If this happens, stop taking Cenomar and contact your doctor immediately.
- You may experience symptoms of neuropathy such as pain, burning, tingling, numbness and/or weakness. If this happens, inform your doctor immediately prior to continuing treatment with Cenomar.
- You may experience mental health problems even when taking quinolone antibiotics, including Cenomar, for the first time. In very rare cases depression or mental health problems have led to suicidal thoughts and self-injurious behaviour such as suicide attempts (see section 4. Possible side effects). If you develop such reactions, stop taking Cenomar and inform your doctor immediately.
- You may develop diarrhoea whilst or after taking antibiotics including Cenomar. If this becomes severe or persistent or you notice that your stool contains blood or mucus you should stop taking Cenomar immediately and consult your doctor. You should not take medicines that stop or slow down bowel movement.
- Cenomar may cause pain and inflammation of your tendons, even within 48 hours of starting treatment and up to several months after discontinuing Cenomar therapy. The risk of inflammation and rupture of tendons is increased if you are elderly or if you are also taking corticosteroids. At the first sign of any pain or inflammation you should stop taking Cenomar, rest the affected limb(s) and consult your doctor immediately. Avoid any unnecessary exercise, as this might increase the risk of a tendon rupture (see sections Do not take Cenomar... and 4. Possible side effects).
- If you are elderly and have kidney problems make sure that you drink plenty whilst taking Cenomar. If you get dehydrated this may increase the risk of kidney failure.
- If your eyesight becomes impaired or if your eyes seem to be otherwise affected, consult an eye specialist immediately (see sections Driving and using machines and 4. Possible side effects).
- Fluoroquinolone antibiotics may cause disturbances in blood sugar, including both a decrease in blood sugar below normal levels (hypoglycemia) and an increase in blood sugar above normal levels (hyperglycemia). In patients treated with Cenomar, disturbances in blood sugar occurred predominantly in elderly patients receiving concomitant treatment with oral antidiabetic medicines that lower blood sugar (e. g. sulfonylurea) or with insulin. If you suffer from diabetes, your blood sugar should be carefully monitored (see section 4. Possible side effects).
- Quinolone antibiotics may make your skin become more sensitive to sunlight or UV light. You should avoid prolonged exposure to sunlight or strong sunlight and should not use a sunbed or any other UV lamp while taking Cenomar.
- The efficacy of Cenomar in the treatment of severe burns, infections of deep tissue and diabetic foot infections with osteomyelitis (infections of the bone marrow) has not been established.

Children and adolescents

Do not give this medicine to children and adolescents under the age of 18 because efficacy and safety have not been established for this age group (see section Do not take Cenomar).

Other medicines and Cenomar

Tell your doctor or pharmacist about any other medicines that you are taking, took recently or might take.

For Cenomar, be aware of the following:

- If you are taking Cenomar and other medicines that affect your heart there is an increased risk for altering your heart rhythm. Therefore, do not take Cenomar together with the following medicines:

- medicines that belong to the group of anti-arrhythmics (e.g. quinidine, hydroquinidine, disopyramide, amiodarone, sotalol, dofetilide, ibutilide)
- antipsychotics (e.g. phenothiazines, pimozide, sertindole, haloperidol, sultopride)
- tricyclic antidepressants
- some antimicrobials (e.g. saquinavir, sparfloxacin, intravenous erythromycin, pentamidine, antimalarials particularly halofantrine)
- some antihistamines (e.g. terfenadine, astemizole, mizolastine)
- other medicines (e.g. cisapride, intravenous vincamine, bepridil and diphemanil).
- You must tell your doctor if you are taking other medicines that can lower your blood potassium levels (e.g. some diuretics, some laxatives and enemas [high doses] or corticosteroids [anti-inflammatory drugs], amphotericin B) or cause a slow heart rate because these can also increase the risk of serious heart rhythm disturbances while taking Cenomar.
- Any medicine containing magnesium or aluminium (such as antacids for indigestion), iron, zinc or didanosine or any medicine containing sucralfate (to treat stomach disorders) can reduce the action of Cenomar tablets. Take your Cenomar tablet 6 hours before or after taking the other medicine.
- Taking any medicine containing charcoal at the same time as Cenomar tablets reduces the action of Cenomar. It is recommended that these medicines are not used together.
- If you are currently taking drugs to thin your blood (oral anti-coagulants such as warfarin), it may be necessary for your doctor to monitor your blood clotting time.

Cenomar with food and drink

Cenomar can be taken with or without food (including dairy products).

Pregnancy, breast-feeding and fertility

Do not take Cenomar if you are pregnant or breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Animal studies do not indicate that your fertility will be impaired by taking this medicine.

Driving and using machines

Cenomar may make you feel dizzy or light-headed, you may experience a sudden, transient loss of vision, or you may faint for a short period. If you are affected do not drive or operate machinery.

Cenomar contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, speak to your doctor before taking Cenomar.

3. HOW TO TAKE CENOMAR

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose for adults is one 400mg film-coated tablet once daily.

Cenomar tablets are for oral use. Swallow the tablet whole (to mask the bitter taste) and with plenty of liquid. You can take Cenomar with or without food. Try to take the tablet at approximately the same time each day.

The same dose can be taken by elderly patients, patients with a low bodyweight or in patients with kidney problems.

The time you will take Cenomar for depends on your infection. Unless your doctor tells you otherwise, your treatment will be as follows:

- for sudden worsening (acute exacerbation) of chronic bronchitis 5 - 10 days
- for infection of the lungs (pneumonia) except for pneumonia which starts during a stay in hospital 10 days
- for acute infection of the sinuses (acute bacterial sinusitis) 7 days

- Mild to moderate infections of the female upper genital tract (pelvic inflammatory disease), including infection of the fallopian tubes and infection of the uterus mucous membrane 14 days

When Cenomar film-coated tablets are used to complete a course of therapy started with Cenomar solution for infusion, the recommended durations of use are:

- Infection of the lungs (pneumonia) acquired outside the hospital 7 -14 days

Most patients with pneumonia were switched to oral treatment with Cenomar film-coated tablets within 4 days.

- Infections of the skin and soft tissue 7 -21 days

Most patients with infections of the skin and soft tissue were switched to oral treatment with Cenomar film-coated tablets within 6 days.

It is important that you complete the course of treatment even if you begin to feel better after a few days. If you stop taking Cenomar too soon your infection may not be completely cured and the infection may return or your condition may get worse. The bacteria causing your infection may become resistant to Cenomar.

The recommended dose and duration of treatment should not be exceeded (see section 2. What you need to know before you take Cenomar, Warnings and precautions).

If you take more Cenomar than you should

If you take more than the prescribed one tablet a day, get medical help immediately. Try to take any remaining tablets, the packaging or this leaflet with you to show the doctor or pharmacist what you have taken.

If you forget to take Cenomar

If you forget to take your tablet you should take it as soon as you remember on the same day. If you do not remember on the same day, take your normal dose (one tablet) on the next day. Do not take a double dose to make up for a forgotten dose.

If you are unsure about what to do ask your doctor or pharmacist.

If you stop taking Cenomar

If you stop taking this medicine too soon your infection may not be completely cured. Talk to your doctor if you wish to stop taking your tablets before the end of the course of treatment.

If you have any further questions about this medicine, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. The most serious side effects observed during treatment with Cenomar are listed below:

If you notice

- an abnormal fast heart rhythm (rare side effect)
- that you suddenly start feeling unwell or notice yellowing of the whites of the eyes, dark urine, itching of the skin, a tendency to bleed or disturbances of thought or wakefulness (these can be signs and symptoms of fulminant inflammation of the liver potentially leading to life-threatening liver failure (very rare side effect, fatal cases have been observed))
- alterations of the skin and mucous membranes like painful blisters in the mouth/nose or at the penis/vagina (Stevens-Johnson syndrome or toxic epidermal necrolysis) (very rare side effects, potentially life threatening)
- a severe, sudden generalised allergic reaction incl. very rarely a life-threatening shock (e.g. difficulty in breathing, drop of blood pressure, fast pulse) (rare side effect)
- swelling incl. swelling of the airway (rare side effect, potentially life-threatening)
- convulsions (rare side effect)
- troubles associated with the nervous system such as pain, burning, tingling, numbness and/or weakness in extremities (rare side effect)
- depression (in very rare cases leading to self-harm, such as suicidal ideations/thoughts, or suicide attempts) (rare side effect)

- insanity (potentially leading to self-harm, such as suicidal ideations/thoughts, or suicide attempts) (very rare side effect)
- severe diarrhoea containing blood and/or mucus (antibiotic associated colitis incl. pseudomembranous colitis), which in very rare circumstances, may develop into complications that are life-threatening (rare side effects)
- pain and swelling of the tendons (tendonitis) (rare side effect) or a tendon rupture (very rare side effect) stop taking Cenomar and tell your doctor immediately as you may need urgent medical advice.

In addition, if you notice

- transient loss of vision (very rare side effect), contact an eye specialist immediately.

If you have experienced life-threatening irregular heart beat (Torsade de Pointes) or stopping of heart beat while taking Cenomar (very rare side effects), tell your treating doctor immediately that you have taken Cenomar and do not restart the treatment.

A worsening of the symptoms of myasthenia gravis has been observed in very rare cases. If this happens, consult your doctor immediately.

If you suffer from diabetes and you notice that your blood sugar is increased or decreased (rare or very rare side effect), inform your doctor immediately.

If you are elderly with existing kidney problems and you notice decrease in urine output, swelling in your legs, ankles or feet, fatigue, nausea, drowsiness, shortness of breath or confusion (these can be signs and symptoms of kidney failure, a rare side effect), consult your doctor immediately.

Other side effects which have been observed during treatment with Cenomar are listed below by how likely they are:

Common (may affect up to 1 in 10 people)

- nausea
- diarrhoea
- dizziness
- stomach and abdominal ache
- vomiting
- headache
- increase of a special liver enzyme in the blood (transaminases)
- infections caused by resistant bacteria or fungi e.g. oral and vaginal infections caused by Candida
- change of the heart rhythm (ECG) in patients with low blood potassium level

Uncommon (may affect up to 1 in 100 people)

- rash
- stomach upset (indigestion/heartburn)
- changes in taste (in very rare cases loss of taste)
- sleep problems (predominantly sleeplessness)
- increase of a special liver enzyme in the blood (gamma-glutamyl-transferase and/or alkaline phosphatase)
- low number of special white blood cells (leukocytes, neutrophils)
- constipation
- itching
- sensation of dizziness (spinning or falling over)
- sleepiness
- wind
- change of the heart rhythm (ECG)
- impaired liver function (incl. increase of a special liver enzyme in the blood (LDH))
- decreased appetite and food intake
- low white blood cells count
- aches and pains such as back, chest, pelvic and extremities pains
- increase of special blood cells necessary for blood clotting
- sweating
- increased specialised white blood cells (eosinophils)

- anxiety
- feeling unwell (predominantly weakness or tiredness)
- shaking
- joint pain
- palpitations
- irregular and fast heart beat
- difficulty in breathing incl. asthmatic conditions
- increase of a special digestive enzyme in the blood (amylase)
- restlessness / agitation
- tingling sensation (pins and needles) and/or numbness
- skin hives
- widening of blood vessels
- confusion and disorientation
- decrease of special blood cells necessary for blood clotting
- visual disturbances incl. double and blurred vision
- decreased blood clotting
- increased blood lipids (fats)
- low red blood cell count
- muscle pain
- allergic reaction
- increase of bilirubin in the blood
- inflammation of the stomach
- dehydration
- severe heart rhythm abnormalities
- dry skin
- angina pectoris

Rare (may affect up to 1 in 1,000 people)

- muscle twitching
- muscle cramp
- hallucination
- high blood pressure
- swelling (of the hands, feet, ankles, lips, mouth, throat)
- low blood pressure
- kidney impairment (incl. increase in special kidney laboratory test results like urea and creatinine)
- inflammation of the liver
- inflammation of the mouth
- ringing/noise in the ears
- jaundice (yellowing of the whites of the eyes or skin)
- impairment of skin sensation
- abnormal dreams
- disturbed concentration
- difficulty in swallowing
- changes in smell (incl. loss of smell)
- balance disorder and poor co-ordination (due to dizziness)
- partial or total loss of memory
- hearing impairment including deafness (usually reversible)
- increased blood uric acid
- emotional instability
- impaired speech
- fainting
- muscle weakness

Very rare (may affect up to 1 in 10,000 people)

- inflammation of joints

- abnormal heart rhythms
- increase of skin sensitivity
- a feeling of self-detachment (not being yourself)
- increased blood clotting
- muscle rigidity
- significant decrease of special white blood cells (agranulocytosis)

Also, there have been very rare cases of the following side effects reported following treatment with other quinolone antibiotics, which might possibly also occur during treatment with Cenomar:

- Increased blood sodium levels
- Increased blood calcium levels
- A special type of reduced red blood cell count (haemolytic anaemia)
- Muscle reactions with muscle cell damage
- Increased sensitivity of the skin to sunlight or UV light.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. By reporting side effects, you can help provide more information on the safety of this medicine.

5. HOW TO STORE CENOMAR

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date stated on the blister and on the carton. The expiry date refers to the last day of that month.

Do not store above 25°C.

Store in the original package in order to protect from moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

This leaflet was last revised in June 2015